



A Program of Wheeler Clinic

800.232.4424 (Voice/TTY) 860.793.9813 (Fax)
www.ctclearinghouse.org



SUICIDE

Frequently Asked Questions

A Library and Resource Center on Alcohol, Tobacco, Other Drugs, Mental Health and Wellness

What should you do if someone tells you they are thinking about suicide?

If someone tells you they are thinking about suicide, you should take their distress seriously, listen nonjudgmentally, and help them get to a professional for evaluation and treatment. People consider suicide when they are hopeless and unable to see alternative solutions to problems. Suicidal behavior is most often related to a mental disorder (depression) or to alcohol or other substance abuse. Suicidal behavior is also more likely to occur when people experience stressful events (major losses, incarceration). If someone is in imminent danger of harming himself or herself, do not leave the person alone. You may need to take emergency steps to get help, such as calling 911. When someone is in a suicidal crisis, it is important to limit access to firearms or other lethal means of committing suicide.

What are the most common methods of suicide?

Firearms are the most commonly used method of suicide for men and women, accounting for 60 percent of all suicides. Nearly 80 percent of all firearm suicides are committed by white males. The second most common method for men is hanging; for women, the second most common method is self-poisoning including drug overdose. The presence of a firearm in the home has been found to be an independent, additional risk factor for suicide. Thus, when a family member or health care provider is faced with an individual at risk for suicide, they should make sure that firearms are removed from the home.

Why do men commit suicide more often than women do?

More than four times as many men as women die by suicide; but women attempt suicide more often during their lives than do men, and women report higher rates of depression. Men and women use different suicide methods. Women in all countries are more likely to ingest poisons than men. In countries where the poisons are highly lethal and/or where treatment resources scarce, rescue is rare and hence female suicides outnumber males.

Who is at highest risk for suicide in the U.S.?

There is a common perception that suicide rates are highest among the young. However, it is the elderly, particularly older white males that have the highest rates. And among white males 65 and older, risk goes up with age. White men 85 and older have a suicide rate that is six times that of the overall national rate. Some older persons are less likely to survive attempts because they are less likely to recuperate. Over 70 percent of older suicide victims have been to their primary care physician within the month of their death, many did not tell their doctors they were depressed nor did the doctor detect it. This has led to research efforts to determine how to best improve physicians' abilities to detect and treat depression in older adults.

What biological factors increase risk for suicide?

Researchers believe that both depression and suicidal behavior can be linked to decreased serotonin in the brain. Low levels of a serotonin metabolite, 5-HIAA, have been detected in cerebral spinal fluid in persons who have attempted suicide, as well as by postmortem studies examining certain brain regions of suicide victims. One of the goals of understanding the biology of suicidal behavior is to improve treatments. Scientists have learned that serotonin receptors in the brain increase their activity in persons with major depression and suicidality, which explains why medications that desensitize or down-regulate these receptors (such as the serotonin reuptake inhibitors, or SSRIs) have been found effective in treating depression. Currently, studies are underway to examine to what extent medications like SSRIs can reduce suicidal behavior.

Can the risk for suicide be inherited?

There is growing evidence that familial and genetic factors contribute to the risk for suicidal behavior. Major psychiatric illnesses, including bipolar disorder, major depression, schizophrenia, alcoholism and substance abuse, and certain personality disorders, which run in families, increase the risk for suicidal behavior. This does not mean that suicidal behavior is inevitable for individuals with this family history; it

(continued)

simply means that such persons may be more vulnerable and should take steps to reduce their risk, such as getting evaluation and treatment at the first sign of mental illness.

Does depression increase the risk for suicide?

Although the majority of people who have depression do not die by suicide, having major depression does increase suicide risk compared to people without depression. The risk of death by suicide may, in part, be related to the severity of the depression. There are also dramatic gender differences in lifetime risk of suicide in depression. Whereas about 7 percent of men with a lifetime history of depression will die by suicide, only 1 percent of women with a lifetime history of depression will die by suicide.

Another way about thinking of suicide risk and depression is to examine the lives of people who have died by suicide and see what proportion of them were depressed. From that perspective, it is estimated that about 60 percent of people who commit suicide have had a mood disorder (e.g., major depression, bipolar disorder, dysthymia). Younger persons who kill themselves often have a substance abuse disorder in addition to being depressed.

Does alcohol and other drug abuse increase the risk for suicide?

A number of recent national surveys have helped shed light on the relationship between alcohol and other drug use and suicidal behavior. A review of minimum-age drinking laws and suicides among youths age 18 to 20 found that lower minimum-age drinking laws was associated with higher youth suicide rates. In a large study following adults who drink alcohol, suicide ideation was reported among persons with depression. In another survey, persons who reported that they had made a suicide attempt during their lifetime were more likely to have had a depressive disorder, and many also had an alcohol and/or substance abuse disorder. In a study of all nontraffic injury deaths associated with alcohol intoxication, over 20 percent were suicides.

In studies that examine risk factors among people who have completed suicide, substance use and abuse occurs more frequently among youth and adults, compared to older persons. For particular groups at risk, such as American Indians and Alaskan Natives, depression and alcohol use and abuse are the most common risk factors for completed suicide. Alcohol and substance abuse problems contribute to suicidal behavior in several ways. Persons who are dependent on substances often have a number of other risk factors for suicide. In addition to being depressed, they are also likely to have social and financial problems. Substance use and abuse can be common among persons prone to be impulsive, and among persons who engage in many types of high risk behaviors that result in self-harm. Fortunately,

there are a number of effective prevention efforts that reduce risk for substance abuse in youth, and there are effective treatments for alcohol and substance use problems. Researchers are currently testing treatments specifically for persons with substance abuse problems who are also suicidal, or have attempted suicide in the past.

What does "suicide contagion" mean, and what can be done to prevent it?

Suicide contagion is the exposure to suicide or suicidal behaviors within one's family, one's peer group, or through media reports of suicide and can result in an increase in suicide and suicidal behaviors. Direct and indirect exposure to suicidal behavior has been shown to precede an increase in suicidal behavior in persons at risk for suicide, especially in adolescents and young adults.

The risk for suicide contagion as a result of media reporting can be minimized by factual and concise media reports of suicide. Reports of suicide should not be repetitive, as prolonged exposure can increase the likelihood of suicide contagion. Suicide is the result of many complex factors; therefore media coverage should not report oversimplified explanations such as recent negative life events or acute stressors. Reports should not divulge detailed descriptions of the method used to avoid possible duplication. Reports should not glorify the victim and should not imply that suicide was effective in achieving a personal goal such as gaining media attention. In addition, information such as hotlines or emergency contacts should be provided for those at risk for suicide.

Following exposure to suicide or suicidal behaviors within one's family or peer group, suicide risk can be minimized by having family members, friends, peers, and colleagues of the victim evaluated by a mental health professional. Persons deemed at risk for suicide should then be referred for additional mental health services.

Is it possible to predict suicide?

At the current time there is no definitive measure to predict suicide or suicidal behavior. Researchers have identified factors that place individuals at higher risk for suicide, but very few persons with these risk factors will actually commit suicide. Risk factors include mental illness, substance abuse, previous suicide attempts, family history of suicide, history of being sexually abused, and impulsive or aggressive tendencies. Suicide is a relatively rare event and it is therefore difficult to predict which persons with these risk factors will ultimately commit suicide.

National Institute of Mental Health (NIMH)
<http://www.nimh.nih.gov/SuicidePrevention/suicidefaq.cfm>